

AD-A167 105

SYNTHESIS OF SOME ARSINO GALLANES AND THE NOVEL
REARRANGEMENT OF A DIMERIC. (U) DUKE UNIV DURHAM NC
DEPT OF CHEMISTRY C G PITT ET AL. APR 86 DU/DC/TR-83
N00014-83-K-0572


1/1

UNCLASSIFIED

F/G 7/3

NL





MICROCOPY

CHART

AD-A167 105

OFFICE OF NAVAL RESEARCH

Contract N00014-83-K0572

Task No. NR 053-841

TECHNICAL REPORT NO. DU/DC/TR-03

Synthesis of Some Arsinogallanes and the Novel
Rearrangement of a Dimeric Bisarsinogallane,
Bis{bis[bis(trimethylsilylmethyl)arsino]chlorogallane}

by

C. G. Pitt, A. P. Purdy, K. T. Higa, and R. L. Wells

Prepared for Publication

in

Organometallics

Duke University
Department of Chemistry
Durham, NC 27706

April, 1986

Reproduction in whole or in part is
permitted for any purpose of the United States Government

This document has been approved for public
release and sale; its distribution is unlimited

DTIC
ELECTE
APR 18 1986
S D

DTIC FILE COPY

36 4 17 037

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER DU/DC/TR-03	2. GOVT ACCESSION NO. AD-A167105	3. RECIPIENT'S CAT. NO. NUMBER
4. TITLE (and Subtitle) Synthesis of Some Arsinogallanes and the Novel Rearrangement of a Dimeric Bisarsinogallane, Bis{bis[bis(trimethylsilylmethyl)arsino]chloro- gallane}	5. TYPE OF REPORT & PERIOD COVERED Technical Report 1986	
6. AUTHOR(s) C.G. Pitt, A.P. Purdy, K.T. Higa, and R.L. Wells	7. PERFORMING ORG. REPORT NUMBER TR-03	
8. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Chemistry Duke University Durham, NC 27706	9. CONTRACT OR GRANT NUMBER(s) N00014-83-K0572	
10. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research	11. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS NR 053-841	
12. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)	13. REPORT DATE April 11, 1986	
	14. NUMBER OF PAGES 16	
	15. SECURITY CLASS. (of this report) Unclassified	
16. DISTRIBUTION STATEMENT (of this report) Approved for public release: distribution unlimited		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from report) Prepared for publication in <u>Organometallics</u>		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Organogallium-arsenic Dimeric bisarsinogallane Dynamic NMR Fluxional molecule		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) See attached		

DD FORM 1473

EDITION OF 1 NOV 65 IS OBSOLETE

1. N. 0102-15-010-5001

Unclassified




SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

**Synthesis of Some Arsinogallanes and the Novel
Rearrangement of a Dimeric Bisarsinogallane,
Bis(bis[bis(trimethylsilylmethyl)arsino]chlorogallane)**

Colin G. Pitt*, Andrew P. Purdy, Kelvin T. Higa, and Richard L. Wells*

Department of Chemistry, Paul M. Gross Chemical Laboratory, Duke University
Durham, North Carolina 27706

Abstract: Silylarsines have been used to synthesize arsinogallanes according to the following equations.



Dynamic NMR spectroscopy studies of one of these, $\{[(Me_3SiCH_2)_2As]_2GaCl\}_2$, show it has novel fluxional properties. Changes of the 1H and ^{13}C NMR spectra were reconciled by rapid exchange of the endo- and exocyclic

150-4-16
 Thermal conductivity (ΔG) + 01-
 bis(trimethylsilylmethyl)arsino groups, and spectra simulated on this basis
 were in excellent agreement with the experimental data. ΔG^\ddagger of 17.0 ± 0.0
 kcal/mol for the ^{13}C exchange was obtained by line shape analysis, and was in
 agreement with the estimate derived from the coalescence temperature. The ΔH
 exchange process between 22 and 57 $^\circ\text{C}$ could be simulated using the same
 kinetic parameters, without invoking pyramidal inversion of the exocyclic
 arsenics.

Accession For	
NTIS CRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution /	
Availability Codes	
Dist	Avail and/or Special
A-1	



**Synthesis of Some Arsinogallanes and the Novel
Rearrangement of a Dimeric Bisarsinogallane,
Bis{bis[bis(trimethylsilylmethyl)arsino]chlorogallane}**

Colin G. Pitt*, Andrew P. Purdy, Kelvin T. Higs, and Richard L. Wells*

Department of Chemistry, Paul M. Gross Chemical Laboratory, Duke University
Durham, North Carolina 27706

Summary: Silylarsines, $R_2AsSiMe_3$ ($R = Me_3SiCH_2$ and Mesityl), have been used to synthesize arsinogallanes. Dynamic NMR spectroscopy studies of one of these, $\{[(Me_3SiCH_2)_2As]_2GaCl\}_2$, show it has novel fluxional properties.

Despite the importance of gallium arsenide in emerging semiconductor technology¹, the chemistry of this elemental combination had, until recently², been restricted to the studies of Coates *et al.*³ in the 1960's. They prepared two monoarsinogallanes by cleavage of one Ga-C bond in Me_3Ga with secondary arsines, resulting in the elimination of methane (Eq. 1). In our hands, this






$R = Me; n = 3$

$R = Ph; n = 2$

method became impractical, and eventually failed, as the steric bulk of the substituents was increased;⁴ it was also found to be limited to the cleavage of a single Ga-C bond, and bis- and trisarsinogallanes, $[(R_2As)_{3-x}GaR_x]_n$

(x = 0, 1), were inaccessible by this route⁵. A number of reports of the synthesis of main group compounds by metathetical elimination of trimethylchlorosilane⁶ prompted us to evaluate this route to arsinogallanes. Here we report its successful application (Eqs. 2-4), as well as the novel fluxional properties of a dimeric bisarsinogallane.



Silylarsines⁷ are readily available from lithium arsenides and trimethylchlorosilane, and their reaction with trichlorogallane proceeds stepwise in hydrocarbon solvents⁸. The only by-product, trimethylchlorosilane, is easily removed in vacuo, a considerable advantage when handling these air and moisture sensitive materials. This method did not yield [(Me₃SiCH₂)₂As]₃Ga; however, it did afford the sterically hindered trisarsinogallane (Mes₂As)₃Ga (Eq. 4) which was first prepared by us from lithium dimesitylarsenide and trichlorogallane and shown to be monomeric by single-crystal x-ray analysis^{2d}. Lack of suitable crystals and the very low

solubility of the mono- and bisarsinogallanes containing mesityl groups (Eqs. 2 and 3), also prepared from lithium dimesitylarsenide and trichlorogallane)^{2d}, prohibited determination of their degree of oligomerization.

The bisarsinogallane $[(\text{Me}_3\text{SiCH}_2)_2\text{As}]_2\text{GaCl}$ was shown by cryoscopic measurements to exist predominantly as the dimer (Eq. 3) in benzene. Its ^1H (300 MHz) NMR spectrum in toluene at 22 °C, reproduced in Figure 1a, consisted of singlets at δ 0.25 and 0.27 (72H, Me_3Si), and 1.65 (8H, endo- CH_2As), and doublets (AB quartet, $^2J_{\text{HH}} = 13.5$ Hz) at 0.94 and 1.76 (8H, exo- CH_2As). The ^{13}C NMR spectrum at 21 °C consisted of four singlets at δ 0.05 and 1.15 (Me_3Si), 7.87 and 8.78 (SiCH_2As). Only a single isomer was detected, assigned the trans structure on steric grounds. The ^1H NMR spectrum was measured using a range of concentrations (5.9-111 mmolar) in an attempt to detect equilibria with monomeric or other oligomeric species; no significant changes in peak intensities of the dimer were observed. However, the relative intensities of the peaks due to trace impurities, $(\text{Me}_3\text{SiCH}_2)_2\text{AsH}$ and $(\text{Me}_3\text{SiCH}_2)_2\text{AsAs}(\text{CH}_2\text{SiMe}_3)_2$, resulting from slight hydrolysis and decomposition of the dimer, respectively⁹, increased somewhat with decreasing concentrations.

On raising the temperature from 22 °C to 97 °C, the ^1H AB quartet collapsed to a broadened doublet which then approached coalescence with the singlet. Likewise, the ^{13}C methyl peaks broadened and coalesced at 78 °C (Fig. 1b). This spectroscopic behavior was reversible, with the exception of the formation of a small amount of the above diarsine. These changes of ^1H and ^{13}C spectra, when coupled with the dimeric molecular weight and the absence of other species in the spectra, can only be reconciled by rapid exchange of the

endo- and exocyclic bis(trimethylsilylmethyl)arsino groups. Spectra simulated on this basis were in excellent agreement with the experimental data. A ΔG^\ddagger of 17.0 ± 0.0 kcal/mol for the ^{13}C exchange was obtained by line shape analysis¹⁰, and was in agreement with the estimate derived from the coalescence temperature¹¹. The ^1H exchange process between 22 and 57 °C could be simulated using the same kinetic parameters, without invoking pyramidal inversion of the exocyclic arsenics. Evidently, the latter is not a significant process in this temperature range, despite the fact that the electropositive gallium might be expected to reduce the inversion barrier.

The equivalency of the bis(trimethylsilylmethyl)arsino groups can be achieved by associative or dissociative mechanisms. The associative mechanism is intuitively more attractive, involving an intermediate in which one or both gallium atoms assume a five-coordinate square pyramidal configuration (Fig. 2). There is ample precedent for five-coordinate gallium.¹² The dissociative mechanism requires cleavage of one endocyclic Ga-As bond, then reclosure of the ring, in a trans-cis-trans equilibrium process. Dissociation into two monomers is considered energetically improbable.

Acknowledgments. The financial support of this work by the Office of Naval Research is gratefully acknowledged. The technical contributions of R. Ernest and C. Foley (Duke University), and T. Caves (North Carolina State University) are similarly recognized.

References and Notes

- (1) Keyes, R.W. Science 1985, 230, 138.
- (2) (a) Wells, R.L.; Purdy, A.P.; McPhail A.T.; Pitt, C.G. Abstracts of Papers, 189th National Meeting of the American Chemical Society, Miami Beach, FL, April 1985; American Chemical Society, Washington, D.C., 1985, INOR 26. (b) Wells, R.L.; Purdy, A.P.; McPhail, A.T.; Pitt, C.G. J. Chem. Soc., Chem. Commun. 1986, in press. (c) Wells, R.L.; Purdy A.P.; McPhail, A.T.; Pitt, C.G. J. Organomet. Chem., submitted. (d) Pitt, C.G.; Higa K.T.; McPhail A.T.; Wells, R.L. Inorg. Chem., submitted.
- (3) (a) Coates, G.E.; Graham, J.J. J. Chem. Soc. 1963, 233. (b) Beachley, O.T.; Coates, G.E. J. Chem. Soc. 1965, 3241.
- (4) For example, $[(\text{Me}_3\text{SiCH}_2)_2\text{AsGaPh}_2]_2$ was prepared by this method (ref. 2c); however, heating Ph_2AsH and $(\text{Me}_3\text{SiCH}_2)_3\text{Ga}$ (1:1 molar ratio) at 60°C for 2 days, afforded impure $[(\text{Ph}_2\text{AsGa}(\text{CH}_2\text{SiMe}_3)_2)_n]$, $\text{Ph}_2\text{AsAsPh}_2$, Me_4Si , H_2 , and unreacted $(\text{Me}_3\text{SiCH}_2)_3\text{Ga}$; no evidence for the desired product was obtained when $(\text{Me}_3\text{SiCH}_2)_2\text{AsH}$ and $(\text{Me}_3\text{SiCH}_2)_3\text{Ga}$ (1:1 molar ratio) were heated at $100\text{--}120^\circ\text{C}$ for 1 month.
- (5) For example, when Ph_2AsH and Me_3Ga (2:1 molar ratio) were heated at $80\text{--}100^\circ\text{C}$ for 2 days, no evidence for any product other than CH_4 , $(\text{Ph}_2\text{AsGaMe}_2)_n$, and unchanged Ph_2AsH was obtained.
- (6) See for example: (a) Abel, E.W.; Armitage, D.A.; Willey, G.R. J. Chem. Soc. 1965, 57. (b) Wells, R.L.; Collins, A.L. Inorg. Chem. 1966, 5, 1327. (c) Abel, E.W.; Illingsworth, S.M. J. Chem. Soc. (A) 1969, 1094. (d) Harman, J.S.; McCartney, M.E.; Sharp, D.W.A. J. Chem. Soc. (A) 1971, 1547. (e) Goetze, R.; Noeth, H. Z. Naturforsch. 1975, 30b, 875. (f) Nutt, W.R.;

Stimson, R. E.; Leopold, M.F.; Rubin, B.H. Inorg. Chem. 1982, 21, 1909. (g)
Hoffmann, G.G. J. Organomet. Chem. 1984, 273, 187.

(7) See for example: (a) Abel, E.W.; Honigschmidt-Grossich, R.; Illingworth, S. M. J. Chem. Soc. (A) 1968, 2623. (b) Abel, E. W; Crow, J.P. J. Organomet. Chem. 1969, 17, 337. (c) Mislow, K.; Senkler, G.H.; Cook, R.J.; Casey, J.P.; Baechler, R. D. J. Am. Chem. Soc. 1972, 94, 2859. The new silylarsines used in this work were prepared under strict inert atmospheric conditions as follows. $(\text{Me}_3\text{SiCH}_2)_2\text{AsSiMe}_3$. A mixture of $(\text{Me}_3\text{SiCH}_2)_2\text{AsH}^{2c}$ (6.0 g, 24 mmol) and *n*-BuLi (10 mL, 2.4 M in hexane) was stirred at 60 °C for 1 day under nitrogen pressure and then cooled to 0 °C, before adding dropwise excess Me_3SiCl (4 mL in 20 mL pentane). After refluxing for 0.5 h, addition of 10 mL of Et_2O , and standing at room temperature for 1 day, the mixture was filtered; following solvent removal, distillation afforded the silylarsine as a colorless liquid (5.8 g, 75% yield), bp 40 °C/0.04-0.35 Torr. Anal. Calcd (Found) for $\text{C}_{11}\text{H}_{31}\text{AsSi}_3$: C, 40.96 (40.83); H, 9.69 (9.82). NMR: ^1H (80 MHz) (C_6D_6) δ 0.80 and 0.46 [AB pattern ($^2J_{\text{HH}} = 13.6$ Hz) 4H, CH_2], 0.18 (s, 9H, AsSiMe_3), 0.15 (s, 18H, CSiMe_3); ^{13}C (22.5 MHz) (C_6D_6) δ 5.7 (s, CH_2), -1.5 (s, AsSiMe_3), 0.37 (s, CSiMe_3). $\text{Mes}_2\text{AsSiMe}_3$. (a) Mesitylmagnesium bromide (274 mmol) in THF (200 mL) was added over a 45 min period, with stirring, to AsCl_3 (25.0 g, 138 mmol) in THF; after 12 h, degassed water (125 mL) and conc HCl (60 mL) were added, and the mixture was stirred for 1 day. Following extraction with benzene, the volatiles were removed by distillation leaving crude, solid Mes_2AsCl (45.1 g, 94% yield). NMR: ^1H (80 MHz) (C_6D_6) δ 6.60 (s, 4H, ring CH), 2.35 (s, 12H, Me), 2.02 (s, 6H Me). (b) The crude Mes_2AsCl was dissolved in THF (500 mL) and LiAlH_4 (10.0 g, 263 mmole) was added. After stirring the solution for 46 h, addition of water (220 mL), extraction with

benzene, and removal of volatiles, distillation afforded Mes_2AsH as a colorless liquid (14.6 g, 58% yield), bp 145–150 °C/0.01 Torr. Anal. Calcd (Found) for $\text{C}_{18}\text{H}_{23}\text{As}$: C, 68.79 (68.66); H, 7.38 (7.55). NMR: ^1H (80 MHz) (C_6D_6) δ 6.70 (s, 4H ring CH), 4.83 (s, 1H, AsH), 2.27 (s, 12H, Me), 2.07 (s, 6H, Me). IR: (Nujol) 2130 cm^{-1} (s, AsH). (c) A mixture of Mes_2AsH (14.62 g, 46.42 mmol), $n\text{-BuLi}$ (20 mL, 2.4 M in hexane), and ligroin (150 mL) was heated at 50 °C for 20 min. Filtration gave solid, yellow Mes_2AsLi which was dried under vacuum (14.41 g, 96.7% yield). Anal. Calcd (Found) for $\text{C}_{18}\text{H}_{22}\text{AsLi}$: C, 67.51 (67.15); H, 7.82 (7.95). NMR: ^1H (80 MHz) (C_6D_6 + 1 drop THF) δ 6.93 (s, 4H, ring CH), 2.61 (s, 12H, Me) 2.29 (s, 6H, Me). (d) A mixture of Mes_2AsLi (3.79 g, 11.8 mmol), Me_3SiCl (10 mL), and ligroin (100 mL) was heated to reflux, stirred at room temperature for 12 h, and filtered. The filtrate was concentrated in vacuo to an oil which eventually crystallized as colorless $\text{Mes}_2\text{AsSiMe}_3$ (4.46 g, 98% yield), mp 62–66 °C Anal. Calcd (Found) for $\text{C}_{21}\text{H}_{31}\text{AsSi}$: C, 65.26 (65.53); H, 8.08 (8.24). NMR: ^1H (80MHz) (C_6D_6) δ 6.74 (s, 4H, ring CH), 2.31 (s, 12H, Me), 2.10 (s, 6H, Me), 0.28 (s, 9H, SiMe_3).

(8) The arsinogallanes were prepared under strict inert atmospheric conditions as follows. $[(\text{Me}_3\text{SiCH}_2)_2\text{AsGaCl}_2]_3$. A solution of $(\text{Me}_3\text{SiCH}_2)_2\text{AsSiMe}_3$ (0.5 g, 1.55 mmol) in pentane (6 mL) was added to a solution of GaCl_3 (0.27 g, 1.55 mmol) in pentane (6 mL). After several min, a precipitate formed. Removal of the liquid phase, washing with pentane, and drying in vacuo gave the arsinogallane as a white solid (0.40 g, 67% yield), mp 170–184 °C (decomp.). Anal. Calcd (Found) for $\text{C}_{24}\text{H}_{66}\text{As}_3\text{Cl}_6\text{Ga}_3\text{Si}_6$: C, 24.64 (24.82); H, 5.68 (5.97); Cl, 18.18 (17.84); mol wt (cryoscopic in benzene), 1170 (1070 80); NMR spectrum shows one species. NMR: ^1H (250 MHz) (C_6D_6) δ 1.78 (s, 12H, CH_2), 0.23 (s, 54H, SiMe_3); ^{13}C (22.5 MHz) (C_6D_6) δ 7.4 (s, CH_2)

1.2 (s, SiMe₃). $\{[(\text{Me}_3\text{SiCH}_2)_2\text{As}]_2\text{GaCl}\}_2$. A solution of $(\text{Me}_3\text{SiCH}_2)_2\text{AsSiMe}_3$ (1.0 g, 3.10 mmol) in pentane (10 mL) was added to a solution of GaCl₃ (0.27 g, 1.55 mmol) in benzene (10 mL) and, after 1 day, the mixture was brought to a boil. Removal of the volatiles under vacuum gave a white solid that on slow recrystallization from ligroin (bp 95-110) afforded the bisarsinogallane as colorless crystals that were dried in vacuo (0.67 g, 80% yield), mp 101-128 °C (decomp.). Anal. Calcd (Found) for C₃₂H₈₈As₄Cl₂Ga₂Si₈: C, 31.82 (31.66); H, 7.34 (7.23); mol wt (cryoscopic in cyclohexane), 1207 (1100 ± 55).

$(\text{Mes}_2\text{AsGaCl}_2)_n$.^{2d} A solution of Mes₂AsSiMe₃ (0.39 g, 1.0 mmol) in pentane (3 mL) was added to a solution of GaCl₃ (0.24 g, 1.36 mmol) in pentane (5 mL) and, after 5 min, the volatiles were removed. Washing the remaining solid with ligroin and drying in vacuo gave the arsinogallane as a white solid (0.46 g, 100% yield), mp 145-146 °C (decomp.). Anal.^{2d} Calcd (Found) for C₁₈H₂₂AsCl₂Ga: C, 47.63 (47.91); H, 4.89 (5.07). NMR: ¹H (80 MHz) (C₆D₆ + 2 drops THF) δ 6.69 (s, 4H, ring CH), 2.45 (s, 12H, Me), 2.06 (s, 6H, Me).

$[(\text{Mes}_2\text{As})_2\text{GaCl}]_n$.^{2d} Addition of a solution of Mes₂AsSiMe₃ (0.82 g, 2.1 mmol) in benzene (10 mL) to a solution of GaCl₃ (0.35 g, 2.0 mmol) in benzene (10 mL) gave a white precipitate of $(\text{Mes}_2\text{AsGaCl}_2)_n$. Addition of more Mes₂AsSiMe₃ (0.75 g, 1.9 mmol) resulted in a yellow-orange solution. Removal of volatiles after 3 h gave a yellow solid that on recrystallization from ligroin/benzene (1:1) afforded the bisarsinogallane as a yellow solid (0.35 g, 24% yield), mp 120-125 °C (decomp.). Anal. Calcd (Found) for C₃₆H₄₄As₂ClGa: C, 59.09 (59.27); H, 6.06 (6.02). NMR: ¹H (80 MHz) (C₆D₆) δ 6.68 (s, 8H, ring CH), 2.31 (s, 24H, Me), 2.06 (s, 12H, Me).

$(\text{Mes}_2\text{As})_3\text{Ga}$.^{2d} Addition of a solution of Mes₂AsSiMe₃ (0.67 g, 1.7 mmol) in benzene (10 mL) to a solution of GaCl₃ (0.15 g, 0.85 mmol) in benzene (10 mL) gave a yellow solution. Removal

of volatiles after 3 days gave a yellow solid which was shown by NMR to be an impure mixture of mono- and bisarsinogallane. Addition of more $\text{Mes}_2\text{AsSiMe}_3$ (0.21 g, 0.54 mmol) dissolved in benzene (10 mL) gave a yellow solid and orange solution. Filtration and removal of volatiles gave an orange oil to which ligroin (10 mL) was added. The resulting mixture was heated and allowed to cool; after 1 day the trisarsinogallane precipitated as an orange solid (0.11 g, 13% yield), mp 130-160 °C (decomp). Anal.^{2d} Calcd (Found) for $\text{C}_{54}\text{H}_{66}\text{As}_3\text{Ga}$: C, 64.24 (64.55); H, 6.59 (6.59). NMR: ^1H (80MHz) (C_6H_6) δ 6.63 (s, 12H, ring CH), 2.32 (s, 36H, Me) 2.05(s, 18H, Me).

(9) Based on NMR, same peaks as in spectra of authentic samples of the arsine^{2c} and the diarsine. The diarsine was prepared by the reduction of $(\text{Me}_3\text{SiCH}_2)_2\text{AsCl}$ ^{2c, 13} with zinc amalgam in refluxing methanol. Anal. Calcd (Found) for $\text{C}_{16}\text{H}_{44}\text{As}_2\text{Si}_4$: C, 38.54 (38.11); H, 8.89 (8.61). NMR: ^1H (80 MHz) (C_6D_6) δ 0.75 and 0.61 [AB pattern ($^2J_{\text{HH}} = 13.6$), 8H, CH_2], 0.12 (s, 36H, SiMe_3); ^{13}C (22.5 MHz) (C_6D_6) δ 10.2 (s, CH_2), 0.5 (s, SiMe_3).

(10) (a) The ^{13}C (75.429 MHz) Me_3Si resonances from spectra at eleven temperatures (304, 309, 314, 319, 325, 330, 349, 350, 351, 352, and 365 °K) were compared with the simulations from the modified DNMR3^{10b} program, using $\Delta\nu = 83.5$ Hz, and a natural line width of 1.0 Hz, and the rate constants (5, 5, 11, 14.5, 29, 35, 150, 170, 185, 195, and 500 sec^{-1}) were obtained. A plot and least-squares analysis of $\ln(k/T)$ versus $1/T$ yielded a slope ($m = -8295 \pm 218$) and an intercept ($b = 22.98 \pm 0.66$). The equations^{10c} $\Delta H^\ddagger = -mR$, $\Delta S^\ddagger = R(b - 23.76)$, and $\Delta G^\ddagger = \Delta H^\ddagger - T\Delta S^\ddagger$ were used to obtain ΔH^\ddagger (16.5 ± 0.4 kcal/mole), ΔS^\ddagger (-1.55 ± 1.3 cal/mol-°K), and ΔG^\ddagger . (b) Binsch, G.; Kleier, D.A. The Computation of Complex Exchange-Broadened NMR Spectra, Program 165, Quantum Chemistry Program Exchange, Indiana University, 1970; revised by

Bushweller, C.H.; Bhat, G.; Letendre, L.J.; Brunelle, J.A.; Bilofsky, H.S.; Ruben, H.; Templeton, D.H.; Zalkin, A. J. Am. Chem. Soc. 1975, 97, 65; and by Caves, T. (North Carolina State University). (c) Sandstrom, J. "Dynamic NMR Spectroscopy"; Academic Press Inc.: London, England, 1982.


(11) $\Delta G_c^\ddagger = 17.0 \pm 0.1$ kcal/mol; obtained (in cal/mol) from the equation $\Delta G_c^\ddagger = T_c [45.67 + 4.58 \log (T_c / \Delta \nu)]$, where $T_c = 351 \pm 1$ °K, and $\Delta \nu = 83.3$ 0.2 Hz.


(12) See for example: (a) Pattison, I.; Wade K. J. Chem. Soc. (A) 1968, 2618. (b) Dymock, K.; Palenik, G.J. J. Chem. Soc., Chem. Commun. 1973, 884. (c) Rettig, S.J.; Storr, A.; Trotter, J. Can. J. Chem. 1975, 53, 753. (d) Rettig, S.J.; Storr, A.; Trotter, J. Can. J. Chem. 1976, 54, 1278. (e) McPhail, A.T.; Miller R.W.; Pitt, C.G.; Gupta, G.; Srivastava, S.C. J. Chem. Soc., Dalton Trans. 1976, 1657.

(13) Seyferth, D. U. S. Patent 2964550 1960.


Figure 1. (a) ^1H NMR Spectra and (b) ^{13}C NMR Spectra
of $\{[(\text{Me}_3\text{SiCH}_2)_2\text{As}]_2\text{GaCl}\}_2$.

Figure 2. Exchange of the Endo- and Exocyclic Arsino Groups.





(b)



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 413 800 N. Quincy Street Arlington, Virginia 22217	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 5042 Crane, Indiana 47522	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Washington, D.C. 20360	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401	1	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
DTNSRDC Attn: Dr. G. Bosmajian Applied Chemistry Division Annapolis, Maryland 21401	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 92132	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	1		

END

FILMED

6-86

DTIC